

Promotor L® 47.0

Pienso complementario líquido / Liquid complementary feed

ES **Componentes analíticos:** Proteina bruta: >13,5 %; Grasa bruta: 0 %; Fibra bruta: 0 %; Cenizas brutas: < 4 %; Metionina > 0,02 %; Lisina > 0,7 %; Calcio, Magnesio: 0 %; Sodio 0,097 %. Fósforo 0,070 %. **Aditivos/1000 ml:** Vitaminas, provitaminas y substancias químicamente definidas de efecto análogo: Niacinamida (3a315) 16,25 g; D-Pantenol (3a842) 7,5 g; Vitamina B1 (3a820) 1,75 g; Vitamina B2 (3a826) 2,5 g; Vitamina B6 (3a831) 1,125 g; Biotina (3a880) 0,001 g. **Composición:** Hidrolizado de levadura de cerveza* y Dextrosa. * El hidrolizado de levadura de cerveza aporta: L-Alanina; L-Arginina; L-Ácido Aspártico; L-Fenilalanina; L-Ácido Glutámico; L-Glicina; L-Histidina; L-Isoleucina; L-Leucina; L-Lisina; L-Metionina; L-Prolina; L-Serina; L-Treonina; L-Tirosina; L-Valina; L-Ácido gamma-aminobutírico, L-Ornitina. **Humedad:** 70-85 %. **Instrucciones de uso:** Desequilibrios y deficiencias de aminoácidos o de vitaminas del grupo B. Oral a través del agua de bebida. Renovar el agua diariamente. Aves: 0,5-1 ml por litro de agua de bebida durante 5 días. Conejos: 2 ml por litro de agua de bebida durante 7 días. Resto de especies: 2 ml por cada 10 kg de peso vivo y día, durante un periodo de 4 a 5 días. Conservar en envase cerrado y ambiente fresco. Nº Autorización fabricante: αESP08100025.

EN **Analytical components:** Gross protein: >13.5 %; Gross fat: 0 %; Gross fiber: 0 %; Ash: < 4 %; Methionine: > 0.02 %; Lysine: > 0.7 %; Calcium, Magnesium: 0 %; Sodium 0.097 %; Phosphorus 0.070 %. **Additives/1000 ml:** Vitamins, provitamins and substances defined chemically of similar effect: Niacinamide (3a315) 16.25 g; D-Panthenol (3a842) 7.5 g; Vitamin B1(3a820) 1.75 g; Vitamin B2 (3a826) 2.5 g; Vitamin B6 (3a831) 1.125 g; Biotin (3a880) 0.001 g. **Composition:** Brewers' yeast hydrolysed * and Dextrose. * The brewers' yeast hydrolysed provides: L-Alanine; L-Arginine; L-Aspartic L-Acid; L-Phenylalanine; L-Glutamic acid; L-Glycine; L-Histidine; L-Isoleucine; L-Leucine; L-Lysine; L-Methionine; L-Proline; L-Serine; L-Threonine; L-Tyrosine; L-Valine; L-Gamma Aminobutyric acid, L-Ornithine. **Humidity:** 70 – 85 %. **Instructions for use:** Aminoacid or group B vitamin deficiencies and disorders. Oral through drinking water. Renew water daily. Poultry: 0.5-1 ml per litre of drinking water for 5 days. Rabbits: 2 ml per litre of drinking water for 7 days. Other species: 2 ml per 10 kg of bodyweight and day, for a period of 4-5 days. Keep the container closed and in a cool place. Nº Authorisation: αESP08100025

Ref. 303420-005

Volumen neto / *Net volume*:
Fab. / *Manuf. Date:*
Lote / *Batch*:
Utilizar preferentemente
antes del / *Best-before*:

ER - All rights
ved

LABORATORIOS CALIER, S.A. C/Barcelones, 26 (El Ramassar) Les Franqueses del Vallès, (Barcelona) España / Spain

CALIER